

## Frozen shoulder: A systematic review of therapeutic options

Harpal Singh Uppal, Jonathan Peter Evans, Christopher Smith

Harpal Singh Uppal, Jonathan Peter Evans, Christopher Smith, Shoulder Unit, Princess Elizabeth Orthopaedic Centre, Royal Devon and Exeter Hospital, EX2 5DW Exeter, United Kingdom

**Author contributions:** All authors contributed equally and there are no conflicts of interests for any authors.

**Open-Access:** This article is an open-access article which was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

**Correspondence to:** Christopher Smith, FRCS, Shoulder Unit, Princess Elizabeth Orthopaedic Centre, Royal Devon and Exeter Hospital, Barrack Road, EX2 5DW Exeter, United Kingdom. [christophersmith3@nhs.net](mailto:christophersmith3@nhs.net)  
Telephone: +44-13-92411611

Received: May 28, 2014

Peer-review started: June 18, 2014

First decision: August 14, 2014

Revised: September 14, 2014

Accepted: October 1, 2014

Article in press: October 10, 2014

Published online: March 18, 2015

Shoulder; Arthroscopic capsular release; Arthrographic distension; Physiotherapy; Steroid; Hydrodilatation

© **The Author(s) 2015.** Published by Baishideng Publishing Group Inc. All rights reserved.

**Core tip:** Frozen shoulder is a common disease which causes significant morbidity. Despite over a hundred years of treating this condition the definition, diagnosis, pathology and most efficacious treatments are still largely unclear. This systematic review of current treatments for frozen shoulder reviews the evidence base behind physiotherapy, both oral and intra articular steroid, hydrodilatation, manipulation under anaesthesia and arthroscopic capsular release. Key areas in which future research could be directed are identified, in particular with regard to the increasing role of arthroscopic capsular release as a treatment.

Uppal HS, Evans JP, Smith C. Frozen shoulder: A systematic review of therapeutic options. *World J Orthop* 2015; 6(2): 263-268 Available from: URL: <http://www.wjgnet.com/2218-5836/full/v6/i2/263.htm> DOI: <http://dx.doi.org/10.5312/wjo.v6.i2.263>

### Abstract

Frozen shoulder is a common disease which causes significant morbidity. Despite over a hundred years of treating this condition the definition, diagnosis, pathology and most efficacious treatments are still largely unclear. This systematic review of current treatments for frozen shoulder reviews the evidence base behind physiotherapy, both oral and intra articular steroid, hydrodilatation, manipulation under anaesthesia and arthroscopic capsular release. Key areas in which future research could be directed are identified, in particular with regard to the increasing role of arthroscopic capsular release as a treatment.

**Key words:** Frozen shoulder; Adhesive capsulitis; Bursitis;

### INTRODUCTION

The first recorded description of a frozen shoulder was reported by Duplay<sup>[1]</sup> in 1872 in his description of a "peri-arthritis scapulohumerale", though the term frozen shoulder was first used in 1934 by Codman<sup>[2]</sup>, who described the common features of a slow onset of pain felt near the insertion of the deltoid muscle, inability to sleep on the affected side, and restriction in both active and passive elevation and external rotation, yet with a normal radiological appearance. Many patients present with a painful restriction of shoulder motion due to pain inhibition or due to weakness from rotator cuff tears or neurological deficits which appear to form a separate clinical entity from patients with no underlying cause for their

symptoms. Patients with secondary frozen shoulder with a clearly identifiable painful primary shoulder pathology often have a poorer prognosis<sup>[3]</sup> and often pose the greatest diagnostic challenges, largely due to the heterogenous nature of the primary pathology. Patients with primary idiopathic frozen shoulder, *i.e.*, patients with a painful global restriction of shoulder movement with no other identifiable shoulder pathology form the basis of this review article.

Frozen shoulder is thought to have an incidence of 3%-5% in the general population and up to 20% in those with diabetes<sup>[4]</sup>. Its peak incidence is between the ages of 40 and 60 and is rare outside these age groups and in manual workers<sup>[3]</sup> and is slightly more common in women. In terms of consultations to general practice it is thought that the cumulative incidence of consultations is 2.4/1000/year (95%CI: 1.9-2.9)<sup>[5]</sup>. Bilateral contemporaneous frozen shoulder occurs in 14% of patients whilst up to 20% of patients will develop some form of similar symptoms in the other shoulder<sup>[6]</sup>. Diabetes is the most common associated disease with frozen shoulder and a patient with diabetes has a lifetime risk of 10%-20% of developing this condition<sup>[7,8]</sup>. Patients with frozen shoulder have a higher risk of having some form of prediabetic condition with an abnormal fasting glucose or impaired glucose tolerance test<sup>[8]</sup>.

Frozen shoulder starts with a painful phase which leads to stiffness which suggests that there is an initial inflammatory response which evolves into a fibrotic reaction. There is some evidence of this occurring histologically<sup>[9]</sup> and there are some similarities to the fibrous contractures in Dupuytren's disease<sup>[10]</sup>. Current models indicate that initial active fibroblastic proliferation in the capsule of the shoulder joint is later accompanied by some transformation of fibroblasts to myofibroblasts<sup>[9,10]</sup>. This thus causes an inflammatory contracture of the shoulder reducing the capsular volume and ultimately restricting glenohumeral movements. The initiating factors that cause this pathoanatomy are poorly understood<sup>[3]</sup>. Current approaches consider the key role of matrix metalloproteinases in the construction of the extracellular matrix and in the various cytokines that control collagen deposition. That drugs such as Marimastat (a synthetic matrix metalloproteinase inhibitor) can induce conditions very similar to primary frozen shoulder and Dupuytren's disease<sup>[11]</sup> is evidence that there may be a common aberrant molecular pathway in these disorders.

The biomechanics of frozen shoulder indicate that the primary pathology can be correlated to contractures of individual structures in the capsule. Gerber demonstrated<sup>[12]</sup> with capsulorrhaphy in cadaveric experiments that restriction of the antero-superior capsule (including the rotator interval, superior glenohumeral ligament and coracohumeral ligament) produces restriction of external rotation in the abducted shoulder whilst anteroinferior capsular

restriction produces restriction of external rotation in the abducted shoulder. Posterior capsular restriction reduces internal rotation of the shoulder and may be present in more severe forms of frozen shoulder<sup>[12]</sup>.

This disorder is thus one of the most common musculoskeletal problems seen in orthopaedics<sup>[4]</sup>. However, despite the ubiquity of this condition and the advances in shoulder surgery over the last fourteen decades there are still many unknowns in deciding what the best treatment options are for this condition<sup>[6]</sup>.

## OPERATIVE INTERVENTIONS

### **Arthroscopic capsular release**

Initial recommendations suggested that arthroscopy has no place in the treatment of frozen shoulder<sup>[13]</sup>. However in the present day arthroscopic capsular release has become increasingly commonplace<sup>[3,4,14]</sup>. The technique requires general anaesthesia and an examination under anaesthesia to document the preoperative range of motion. Standard posterior and anterior portals are made, a diagnostic arthroscopy is performed to confirm the diagnosis and a synovectomy of the rotator interval is performed. The capsular release starts with excision of the rotator interval to the under surface of the conjoint tendon, the release is extended inferiorly posterior to the tendon of subscapularis down to the five o'clock position. Some surgeons advocate release of the superior edge of subscapularis<sup>[15]</sup>, though this is highly controversial. The superior release is then extended to reach the long head of biceps and is continued to release the coracohumeral ligament in the plane between the superior glenoid and supraspinatus. If internal rotation of the shoulder is significantly restricted then the camera portal can be reversed to facilitate a posterior capsular release. Some surgeons complete the inferior release with a gentle manipulation but some surgeons advocate a full 360 degree capsulectomy under direct vision whilst accepting the higher risk of iatrogenic injury the axillary nerve<sup>[14]</sup>. A randomised study by Chen *et al*<sup>[16]</sup> demonstrated that not performing any form of inferior release, such as a manipulation at the end of surgery, results in poorer functional outcome and range of motion at three months post intervention, though these differences are not maintained at longer follow up points.

A systematic review was conducted using the following search strategy '{joint capsule release" (MeSH Terms) OR ["joint"(All Fields) AND "capsule" (All Fields) AND "release"(All Fields)] OR "joint capsule release"(All Fields) OR ["capsular"(All Fields) AND "release"(All Fields)] OR "capsular release"(All Fields)} AND {"bursitis"(MeSH Terms) OR "bursitis"(All Fields) OR ["frozen"(All Fields) AND "shoulder"(All Fields)] OR "frozen shoulder"(All Fields)}' in pubmed on May 11<sup>th</sup> 2014. Embase and cochrane databases were also searched with the same search strategy and the references of selected journals were scanned to try to

**Table 1** Reviewed studies investigating arthroscopic capsular release as a treatment for primary frozen shoulder

	Year	Patients	Outcome measure(s)	Outcome score pre intervention (standard deviation or range)	Outcome score post intervention (standard deviation or range)	Complications
Smith <i>et al</i> <sup>[6]</sup>	2014	136	OSS, VAS	19.2 (7.4)	38.1 (8.6)	One portal site superficial infection - treated oral antibiotics
Jerosch <i>et al</i> <sup>[17]</sup>	2012	91	Constant	42 (19-58)	85 (36-100)	One shoulder infection - debridement required
Le Lievre <i>et al</i> <sup>[18]</sup>	2012	43	Likert		All 43 pain free on Likert score at 5-12 yr from surgery	Nil
Waszczykowski <i>et al</i> <sup>[19]</sup>	2010	16	Modified constant score (0-75), ASES	19.3	65.9	Nil
Cinar <i>et al</i> <sup>[20]</sup>	2010	26	Constant, UCLA	30.4 (6.2)	82 (18.2)	Nil
Baums <i>et al</i> <sup>[21]</sup>	2006	30	ASES, VAS, SF36	35 (10-70)	91 (62-96)	One case of delayed healing of portal site (no infection), one haematoma
Klinger <i>et al</i> <sup>[22]</sup>	2001	36	Constant	29 (14-51)	66 (35-91)	Nil
Ogilvie-Harris <i>et al</i> <sup>[23]</sup>	1997	17	ASES	2 patients mild pain, 6 in moderate pain, 8 in severe pain	11 pain free, 4 in mild pain, 1 in moderate pain, 1 in severe pain	Nil
Segmüller <i>et al</i> <sup>[24]</sup>	1995	24	Modified constant score	10/20	18/20	Nil

OSS: Oxford shoulder score; ASES: American shoulder and elbow score; VAS: Visual analogue pain score; UCLA: UCLA shoulder score; SF36: Short form 36.

find more studies.

### Inclusion criteria

Clinical studies investigating arthroscopic capsular release to treat primary idiopathic frozen shoulder; studies in English.

### Exclusion criteria

Review articles; studies investigating arthroscopic capsular release in conjunction with another surgical procedure; studies with less than fifteen participants; Double publication of data.

Studies on patients with secondary frozen shoulder: 76 Studies were identified; 18 articles were shortlisted for further review following application of eligibility criteria on published abstracts.

Closer examination of these studies revealed: 2 studies included data that had been published twice; 4 studies were not available in English; 2 studies reported results on arthroscopic capsular release and subacromial decompression; One study investigating a mixture of primary and secondary frozen shoulder with no separation of data analysis.

Nine studies<sup>[6,17-24]</sup> were eligible for review and the results of the data abstraction are compiled in Table 1. This review includes the treatment of 419 patients with primary frozen shoulder. All studies demonstrated a rapid statistically significant increase in postoperative shoulder function following capsular release. Five studies used the Constant-Murley score as the primary outcome measure. The Constant-Murley score is a commonly used measure of shoulder function which unfortunately has very little formal validation<sup>[25]</sup>. Other outcome measures used with more validation include the oxford shoulder score in Smith *et al*<sup>[6]</sup>, Likert score

in Le Lievre *et al*<sup>[18]</sup> and American shoulder and elbow score in Waszczykowski *et al*<sup>[19]</sup> and Baums *et al*<sup>[21]</sup>. None of the studies included any comparative control groups which forms the largest weakness in the current evidence base behind arthroscopic capsular release. Overall, the evidence reviewed demonstrates that arthroscopic capsular release appears to be a safe and effective treatment that can provide a rapid improvement in patient reported shoulder function.

### Manipulation under anaesthesia

In this technique a general anaesthetic is administered and the shoulder joint capsule is gently stretched by moving the humerus into flexion, abduction and finally (optionally) by moving the adducted humerus into external rotation. Great care must be taken to minimise the lever arm used and to maximise the surface area of the arm to which pressure is applied. The largest risk in this procedure is of iatrogenic damage to the upper limb including, humeral fracture, glenohumeral dislocation, rotator cuff tears, glenoid fractures, brachial plexus injuries, labral tears and haematomas<sup>[14]</sup>. It has been demonstrated in post manipulation arthroscopy<sup>[26]</sup> that the typical appearances are of haemarthrosis and capsular tearing but other lesions often seen include iatrogenic superior labral anterior posterior tears, partial subscapularis ruptures and rupture of the anterior labrum. Manipulation under anaesthesia has been shown to be an efficacious treatment<sup>[27]</sup>. However, the results of manipulation when compared to hydrodilatation<sup>[28]</sup> and steroid injection<sup>[29]</sup> are equivocal at best.

### Non-operative treatments

**Hydrodilatation (arthrographic distension):** This

treatment involves the injection of local anaesthetic into the capsule at a pressure high enough to distend and stretch the joint capsule. This procedure first described by Andren *et al*<sup>[30]</sup> does not need to be performed in the operating theatre but is often associated with poor tolerance due to the painful nature of the distension<sup>[4]</sup>. Buchbinder *et al*<sup>[31]</sup>'s systematic cochrane review of hydrodilatation searched MEDLINE, EMBASE, CINAHL and CENTRAL databases from 1966 till November 2006 for studies investigating hydrodilatation type procedures in the treatment of frozen shoulder. These searches were repeated from November 2006 till May 2014 and a total of 7 extra studies were identified two of which were randomised comparative studies<sup>[28]</sup>.

Buchbinder *et al*<sup>[32]</sup>'s randomised controlled study of 46 patients compared hydrodilatation to placebo and demonstrated a statistically and clinically significant improvement in functional outcome scores (shoulder pain and disability index) to 6 wk following intervention but this was not maintained at follow up points beyond this.

Three studies compared hydrodilatation with steroid to intra articular steroid injection alone<sup>[33-35]</sup>. Gam *et al*<sup>[33]</sup>'s and Corbeil *et al*<sup>[34]</sup>'s studies had weaknesses in study construction especially with regard to randomisation systems, elimination of systematic bias and in sample size calculation. Tveitå *et al*<sup>[35]</sup>'s study on the other hand is a well constructed study which scores highly against the Consort criteria<sup>[36]</sup>. Gam *et al*<sup>[33]</sup>, Corbeil *et al*<sup>[34]</sup> and Tveitå *et al*<sup>[35]</sup> all failed to demonstrate any statistically significant differences in functional outcome compared to steroid injection at any outcome point. Gam *et al*<sup>[33]</sup> did report an increase in the range of shoulder motion of the hydrodilatation group as compared to the steroid group. However, given that range of motion is an unvalidated and poor measure of shoulder function it is difficult to make generalisable recommendations on this evidence. Khan *et al*<sup>[37]</sup> compared hydrodilatation and physiotherapy to physiotherapy alone in 36 patients in this quasi randomised and underpowered study. Khan *et al*<sup>[37]</sup> demonstrated statistically significant improvements in range of motion at eight weeks but no differences in visual analogue pain scores.

Jacobs *et al*<sup>[38]</sup> reported results of a three way randomised study comparing a mixture of low volume local anaesthetic and air, intra articular steroid and local anaesthetic with air and steroid. Though this study claims to be investigating arthrographic distension, the low volumes used (3 mL of air in distension group) mean that the study design does not pass the test of face validity. Given that all comparative studies use twenty to forty millilitres of saline, which is many orders of magnitude less compressible than air, it seems very unlikely that any patients capsule was distended in any meaningful way in this study.

Quraishi *et al*<sup>[28]</sup> reported results of small randomised study comparing hydrodilatation to manipulation under anaesthesia. Though no differences were found in

Constant score at any point up to six months following intervention both groups made a clinically significant improvement following intervention.

The major side effect of hydrodilatation appears to be of pain during the procedure<sup>[32,33,35]</sup> though Gam also reported one instance of stroke which was not thought to be related to the intervention.

This systematic review of hydrodilatation demonstrates that this technique appears to be efficacious but there is no good evidence to suggest any superiority to other treatments. High quality randomised studies comparing hydrodilatation to other common treatments, such as arthroscopic capsular release, are needed.

### **Physiotherapy**

Most patients are initially prescribed a course of physiotherapy prior to referral to a surgeon. The aim behind most regimens is to prevent further reduction in range of motion and eventually to increase the range of motion in the affected shoulder. Passive mobilisation and capsular stretching are two of the most commonly used techniques. Despite the near universal use of physiotherapy as a first line treatment for frozen shoulder there is very little high quality evidence to support its use. Cochrane reviews have demonstrated that the current literature base shows that physiotherapy alone has little to no benefit as compared to control groups<sup>[39]</sup>. There are a number of adjuncts that are often used with physiotherapy including extracorporeal shockwave therapy, electromagnetic stimulation, acupuncture and the use of lasers, none of which have been subjected to investigation with randomised controlled studies<sup>[3]</sup>.

### **Steroid injection**

Steroid injection is another almost ubiquitous intervention in frozen shoulder. Multiple cochrane reviews have noted the eventual location of a blind glenohumeral or subacromial injection is highly variable<sup>[31,40]</sup>. The most recent cochrane review collates the information from 26 very heterogeneous studies<sup>[40]</sup> and concludes that there is at best a small short term benefit to steroid injection alone for frozen shoulder but that the evidence base is poor. The difficulty in extracting the effect of steroid from that of physiotherapy, an intervention with which it is often combined in studies has long been noted<sup>[41]</sup>.

### **Oral steroid**

This treatment is rarely prescribed by surgeons, however to date, five trials have been conducted investigating oral steroid therapy, comparing steroid to placebo<sup>[32,42]</sup>, no treatment<sup>[43]</sup>, intra articular injection<sup>[44]</sup> and in conjunction with manipulation under anaesthesia<sup>[45]</sup>. These trials were reviewed in a systematic cochrane review in 2006<sup>[46]</sup> and showed that there is a mild short term (under 6 wk) benefit to oral steroid therapy but that this is not maintained in the longer term. This small short term benefit must be offset against the well known

side effects and risks of oral steroid therapy.

## CONCLUSION

Frozen shoulder is a common disease which causes significant morbidity. Despite over a hundred years of treating this condition the definition, diagnosis, pathology and most efficacious treatments are still largely unclear. This review of the recent evidence base highlights key areas for future research in particular with regard to the increasing role of arthroscopic capsular release as a treatment. High quality adequately powered randomised controlled trials comparing the most common interventions to a sham procedure would be the ideal way to improve the current evidence base. However these are difficult studies to construct and recruit for. Frozen shoulder can be such an intensely painful condition that in severe cases one could consider that an option of no treatment as part of a control group could be considered to be unethical. Given these real world problems in construction of clinical trials the optimum area to concentrate further research is in comparing treatments like arthroscopic capsular release to hydrodilatation with an adequately powered high quality randomised controlled trial.

## REFERENCES

- 1 Duplay E. De la periarthrite scapulo-humérale et des raideurs de l'épaule qui en sont la conséquence. *Arch Gen Med* 1872; **20**: 513-542
- 2 Codman EA. Tendinitis of the Short Rotators. In: The Shoulder: Rupture of the Supraspinatus Tendon and Other Lesions in or about the Subacromial Bursa. Boston MA: Thomas Todd, 1934
- 3 Robinson CM, Seah KT, Chee YH, Hindle P, Murray IR. Frozen shoulder. *J Bone Joint Surg Br* 2012; **94**: 1-9 [PMID: 22219239 DOI: 10.1302/0301-620X.94B1.27093]
- 4 Manske RC, Prohaska D. Diagnosis and management of adhesive capsulitis. *Curr Rev Musculoskelet Med* 2008; **1**: 180-189 [PMID: 19468904 DOI: 10.1007/s12178-008-9031-6]
- 5 van der Windt DA, Koes BW, de Jong BA, Bouter LM. Shoulder disorders in general practice: incidence, patient characteristics, and management. *Ann Rheum Dis* 1995; **54**: 959-964 [PMID: 8546527 DOI: 10.1136/ard.54.12.959]
- 6 Smith CD, Hamer P, Bunker TD. Arthroscopic capsular release for idiopathic frozen shoulder with intra-articular injection and a controlled manipulation. *Ann R Coll Surg Engl* 2014; **96**: 55-60 [PMID: 24417832 DOI: 10.1308/003588414X13824511650452]
- 7 Lundberg BJ. The frozen shoulder. Clinical and radiographical observations. The effect of manipulation under general anesthesia. Structure and glycosaminoglycan content of the joint capsule. Local bone metabolism. *Acta Orthop Scand Suppl* 1969; **119**: 1-59 [PMID: 4952729]
- 8 Tighe CB, Oakley WS. The prevalence of a diabetic condition and adhesive capsulitis of the shoulder. *South Med J* 2008; **101**: 591-595 [PMID: 18475240 DOI: 10.1097/SMJ.0b013e3181705d39]
- 9 Hand GC, Athanasou NA, Matthews T, Carr AJ. The pathology of frozen shoulder. *J Bone Joint Surg Br* 2007; **89**: 928-932 [PMID: 17673588 DOI: 10.1302/0301-620X.89B7.19097]
- 10 Bunker TD, Anthony PP. The pathology of frozen shoulder. A Dupuytren-like disease. *J Bone Joint Surg Br* 1995; **77**: 677-683 [PMID: 7559688]
- 11 Hutchinson JW, Tierney GM, Parsons SL, Davis TR. Dupuytren's disease and frozen shoulder induced by treatment with a matrix metalloproteinase inhibitor. *J Bone Joint Surg Br* 1998; **80**: 907-908 [PMID: 9768907]
- 12 Gerber C, Werner CM, Macy JC, Jacob HA, Nyffeler RW. Effect of selective capsulorrhaphy on the passive range of motion of the glenohumeral joint. *J Bone Joint Surg Am* 2003; **85-A**: 48-55 [PMID: 12533571]
- 13 Neviasser RJ, Neviasser TJ. The frozen shoulder. Diagnosis and management. *Clin Orthop Relat Res* 1987; **(223)**: 59-64 [PMID: 3652593]
- 14 Hsu JE, Anakwenze OA, Warrender WJ, Abboud JA. Current review of adhesive capsulitis. *J Shoulder Elbow Surg* 2011; **20**: 502-514 [PMID: 21167743 DOI: 10.1016/j.jse.2010.08.023]
- 15 Liem D, Alci S, Dedy N, Steinbeck J, Marquardt B, Möllenhoff G. Clinical and structural results of partial supraspinatus tears treated by subacromial decompression without repair. *Knee Surg Sports Traumatol Arthrosc* 2008; **16**: 967-972 [PMID: 18712359 DOI: 10.1007/s00167-008-0580-4]
- 16 Chen J, Chen S, Li Y, Hua Y, Li H. Is the extended release of the inferior glenohumeral ligament necessary for frozen shoulder? *Arthroscopy* 2010; **26**: 529-535 [PMID: 20362834 DOI: 10.1016/j.arthro.2010.02.020]
- 17 Jerosch J, Nasef NM, Peters O, Mansour AM. Mid-term results following arthroscopic capsular release in patients with primary and secondary adhesive shoulder capsulitis. *Knee Surg Sports Traumatol Arthrosc* 2013; **21**: 1195-1202 [PMID: 22763569 DOI: 10.1007/s00167-012-2124-1]
- 18 Le Lievre HM, Murrell GA. Long-term outcomes after arthroscopic capsular release for idiopathic adhesive capsulitis. *J Bone Joint Surg Am* 2012; **94**: 1208-1216 [PMID: 22760389 DOI: 10.2106/JBJS.J.00952]
- 19 Waszczykowski M, Fabiś J. The results of arthroscopic capsular release in the treatment of frozen shoulder - two-year follow-up. *Ortop Traumatol Rehabil* 2010; **12**: 216-224 [PMID: 20675863]
- 20 Cinar M, Akpinar S, Derincek A, Circi E, Uysal M. Comparison of arthroscopic capsular release in diabetic and idiopathic frozen shoulder patients. *Arch Orthop Trauma Surg* 2010; **130**: 401-406 [PMID: 19471947 DOI: 10.1007/s00402-009-0900-2]
- 21 Baums MH, Spahn G, Nozaki M, Steckel H, Schultz W, Klinger HM. Functional outcome and general health status in patients after arthroscopic release in adhesive capsulitis. *Knee Surg Sports Traumatol Arthrosc* 2007; **15**: 638-644 [PMID: 17031613 DOI: 10.1007/s00167-006-0203-x]
- 22 Klinger HM, Otte S, Baums MH, Haerer T. Early arthroscopic release in refractory shoulder stiffness. *Arch Orthop Trauma Surg* 2002; **122**: 200-203 [PMID: 12029508 DOI: 10.1007/s00402-001-0355-6]
- 23 Ogilvie-Harris DJ, Myerthall S. The diabetic frozen shoulder: arthroscopic release. *Arthroscopy* 1997; **13**: 1-8 [PMID: 9043598 DOI: 10.1016/S0749-8063(97)90203-6]
- 24 Segmüller HE, Taylor DE, Hogan CS, Saies AD, Hayes MG. Arthroscopic treatment of adhesive capsulitis. *J Shoulder Elbow Surg* 1995; **4**: 403-408 [PMID: 8665283 DOI: 10.1016/S1058-2746(05)80030-8]
- 25 Harvie P, Pollard TC, Chennagiri RJ, Carr AJ. The use of outcome scores in surgery of the shoulder. *J Bone Joint Surg Br* 2005; **87**: 151-154 [DOI: 10.1302/0301-620X.87B2.15305]
- 26 Loew M, Heichel TO, Lehner B. Intraarticular lesions in primary frozen shoulder after manipulation under general anesthesia. *J Shoulder Elbow Surg* 2005; **14**: 16-21 [PMID: 15723009 DOI: 10.1016/j.jse.2004.04.004]
- 27 Dodenhoff RM, Levy O, Wilson A, Copeland SA. Manipulation under anesthesia for primary frozen shoulder: effect on early recovery and return to activity. *J Shoulder Elbow Surg* 2000; **9**: 23-26 [PMID: 10717858 DOI: 10.1016/S1058-2746(00)90005-3]
- 28 Quraishi NA, Johnston P, Bayer J, Crowe M, Chakrabarti AJ. Thawing the frozen shoulder. A randomised trial comparing manipulation under anaesthesia with hydrodilatation. *J Bone Joint Surg Br* 2007; **89**: 1197-1200 [PMID: 17905957 DOI: 10.1302/0301-620X.89B9.18863]
- 29 Kivimäki J, Pohjolainen T. Manipulation under anesthesia for frozen shoulder with and without steroid injection. *Arch Phys Med*

- Rehabil* 2001; **82**: 1188-1190 [PMID: 11552189 DOI: 10.1053/apmr.2001.24169]
- 30 **Andren L**, Lundberg BJ. Treatment of rigid shoulders by joint distension during arthrography. *Acta Orthop Scand* 1965; **36**: 45-53 [PMID: 14308098 DOI: 10.3109/17453676508989370]
- 31 **Buchbinder R**, Green S, Youd JM, Johnston RV, Cumpston M. Arthrographic distension for adhesive capsulitis (frozen shoulder). *Cochrane Database Syst Rev* 2008; : CD007005 [PMID: 18254123 DOI: 10.1002/14651858.CD007005]
- 32 **Buchbinder R**, Green S, Forbes A, Hall S, Lawler G. Arthrographic joint distension with saline and steroid improves function and reduces pain in patients with painful stiff shoulder: results of a randomised, double blind, placebo controlled trial. *Ann Rheum Dis* 2004; **63**: 302-309 [PMID: 14962967 DOI: 10.1136/ard.2002.004655]
- 33 **Gam AN**, Schydrowsky P, Rossel I, Remvig L, Jensen EM. Treatment of "frozen shoulder" with distension and glucocorticoid compared with glucocorticoid alone. A randomised controlled trial. *Scand J Rheumatol* 1998; **27**: 425-430 [PMID: 9855212 DOI: 10.1080/030097498442244]
- 34 **Corbeil V**, Dussault RG, Leduc BE, Fleury J. [Adhesive capsulitis of the shoulder: a comparative study of arthrography with intra-articular corticotherapy and with or without capsular distension]. *Can Assoc Radiol J* 1992; **43**: 127-130 [PMID: 1562888]
- 35 **Tveitå EK**, Tariq R, Sesseng S, Juel NG, Bautz-Holter E. Hydrodilatation, corticosteroids and adhesive capsulitis: a randomized controlled trial. *BMC Musculoskelet Disord* 2008; **9**: 53 [PMID: 18423042 DOI: 10.1186/1471-2474-9-53]
- 36 **Altman DG**, Schulz KF, Moher D, Egger M, Davidoff F, Elbourne D, Gøtzsche PC, Lang T. The revised CONSORT statement for reporting randomized trials: explanation and elaboration. *Ann Intern Med* 2001; **134**: 663-694 [PMID: 11304107 DOI: 10.7326/003-4819-134-8-200104170-00012]
- 37 **Khan AA**, Mowla A, Shakoor MA, Rahman MR. Arthrographic distension of the shoulder joint in the management of frozen shoulder. *Mymensingh Med J* 2005; **14**: 67-70 [PMID: 15695959]
- 38 **Jacobs LG**, Barton MA, Wallace WA, Ferrous J, Dunn NA, Bossingham DH. Intra-articular distension and steroids in the management of capsulitis of the shoulder. *BMJ* 1991; **302**: 1498-1501 [PMID: 1855018]
- 39 **Buchbinder R**, Green S, Youd JM. Corticosteroid injections for shoulder pain. *Cochrane Database Syst Rev* 2003; : CD004016 [PMID: 12535501 DOI: 10.1002/14651858.CD004016]
- 40 **Green S**, Buchbinder R, Hetrick S. Physiotherapy interventions for shoulder pain. *Cochrane Database Syst Rev* 2003; : CD004258 [PMID: 12804509 DOI: 10.1002/14651858.CD004258]
- 41 **Buchbinder R**, Youd JM, Green S, Stein A, Forbes A, Harris A, Bennell K, Bell S, Wright WJ. Efficacy and cost-effectiveness of physiotherapy following glenohumeral joint distension for adhesive capsulitis: a randomized trial. *Arthritis Rheum* 2007; **57**: 1027-1037 [PMID: 17665470 DOI: 10.1002/art.22892]
- 42 **Blockey NJ**, Wright JK, Kellgren JH. Oral cortisone therapy in periarthritis of the shoulder; a controlled trial. *Br Med J* 1954; **1**: 1455-1457 [PMID: 13160496]
- 43 **Binder A**, Hazleman BL, Parr G, Roberts S. A controlled study of oral prednisolone in frozen shoulder. *Br J Rheumatol* 1986; **25**: 288-292 [PMID: 3730737 DOI: 10.1093/rheumatology/25.3.288]
- 44 **Widiastuti-Samekto M**, Sianturi GP. Frozen shoulder syndrome: comparison of oral route corticosteroid and intra-articular corticosteroid injection. *Med J Malaysia* 2004; **59**: 312-316 [PMID: 15727375]
- 45 **Kessel L**, Bayley I, Young A. The upper limb: the frozen shoulder. *Br J Hosp Med* 1981; **25**: 334, 336-37, 339 [PMID: 7236953]
- 46 **Buchbinder R**, Green S, Youd JM, Johnston RV. Oral steroids for adhesive capsulitis. *Cochrane Database Syst Rev* 2006; **(4)**: CD006189 [PMID: 17054278 DOI: 10.1002/14651858.CD006189]

**P- Reviewer:** Daglar B, Lin JJ, Swanik C, Zheng N **S- Editor:** Ji FF  
**L- Editor:** A **E- Editor:** Liu SQ

